



Printed Pages : 7

EAS-103

(Following Paper ID and Roll No. to be filled in your Answer Book)	
<b>PAPER ID : 9601</b>	Roll No. <input type="text"/>

**B. Tech.**

(Only for the candidates admitted/Readmitted in the session 2008-09)

**(SEM. I) EXAMINATION, 2008-09**

**MATHEMATICS - I**

*Time : 3 Hours]*

*[Total Marks : 100*

**SECTION - A**

All parts of this question are **compulsory**. 2×10=20

1 (a) For which value of 'b' the rank of the matrix

$$A = \begin{bmatrix} 1 & 5 & 4 \\ 0 & 3 & 2 \\ b & 13 & 10 \end{bmatrix} \text{ is } 2, b = \underline{\hspace{2cm}}$$

(b) Determine the constants *a* and *b* such that

the curl of vector  $\vec{A} = (2xy + 3yz)\hat{i} +$

$(x^2 + axz - 4z^2)\hat{j} - (3xy + byz)\hat{k}$  is zero,

*a* = \_\_\_\_\_, *b* = \_\_\_\_\_.

(c) The *n*<sup>th</sup> derivative  $(y_n)$  of the function

$y = x^2 \sin x$  at  $x = 0$  is \_\_\_\_\_.



(d) With usual notations, match the items on right hand side with those on left hand side for properties of  $\text{Max}^m$  and minimum. :

(i)  $\text{Max}^m$  (p)  $rt - s^2 = 0$

(ii)  $\text{Min}^m$  (q)  $rt - s^2 < 0$

(iii) Saddle point (r)  $rt - s^2 > 0, r > 0$

(iv) Failure case (s)  $rt - s^2 > 0$  and  $r < 0$

(e) Match the items on the right hand side with those on left hand side for the following special functions :  
(Full marks is awarded if all matchings are correct)

(i)  $\beta(p, q)$  (p)  $\left(\frac{1}{2}\right)$

(ii)  $\frac{\sqrt{p} \sqrt{q}}{\sqrt{p+q}}$  (q)  $\int_0^\infty \frac{y^{p-1}}{(1+y)(p+q)} dy$

(iii)  $\sqrt{\pi}$  (r)  $\beta(p, q)$

(iv)  $\frac{\pi}{\sin p\pi}$  (s)  $\sqrt{p} \sqrt{1-p}$

**Indicate True or False for the following statements :**

(f) (i) If  $|A| = 0$ , then at least one eigen value is zero. (True / False)

(ii)  $A^{-1}$  exists iff 0 is an eigen value of  $A$ . (True / False)

(iii) If  $|A| \neq 0$ , then  $A$  is known as singular matrix. (True / False)



- (iv) Two vectors X and Y is said to be orthogonal  $Y, X^T Y = Y^T X \neq 0$ . (True / False)
- (g) (i) The curve  $y^2 = 4ax$  is symmetric about x-axis. (True / False)
- (ii) The curve  $x^3 + y^3 = 3axy$  is symmetric about the line  $y = -x$ . (True / False)
- (iii) The curve  $x^2 + y^2 = a^2$  is symmetric about both the axis x and y. (True / False)
- (iv) The curve  $x^3 - y^3 = 3axy$  is symmetric about the line  $y = x$ . (True / False)

**Pick the correct answer of the choices given below :**

- (h) If  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  is position vector, then value of  $\nabla (\log r)$  is

- (i)  $\frac{\vec{r}}{r}$                       (ii)  $\frac{\vec{r}}{r^2}$
- (iii)  $-\frac{\vec{r}}{r^3}$                       (iv) none of the above.

- (i) The Jacobian  $\frac{\partial(uv)}{\partial(xy)}$  for the function

$u = e^x \sin y, v = (x + \log \sin y)$  is

- (i) 1                      (ii)  $\sin x \sin y - xy \cos x \cos y$
- (iii) 0                      (iv)  $\frac{e^x}{x}$



- (j) The volume of the solid under the surface  $az = x^2 + y^2$  and whose base  $R$  is the circle  $x^2 + y^2 = a^2$  is given as
- (i)  $\pi | 2a$       (ii)  $\pi a^3 | 2$
- (iii)  $\frac{4}{3} \pi a^3$       (iv) None of the above.

### SECTION - B

Attempt any **three** parts of the following :

**10×3=30**

- 2 (a) If  $y = (\sin^{-1} x)^2$  prove that  $y_n(0) = 0$  for  $b$  odd and  $y_n(0) = 2, 2^2, 4^2, 6^2 \dots (n-2)^2$ ,  $n \neq 2$  for  $n$  is even.
- (b) Find the dimension of rectangular box of maximum capacity whose surface area is given when (a) box is open at the top (b) box is closed.
- (c) Find a matrix  $P$  which diagonalizes the matrix  $A = \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix}$ , verify  $P^{-1}AP = D$  where  $D$  is the diagonal matrix.
- (d) Find the area and the mass contained  $m$  the first quadrant enclosed by the curve  $\left(\frac{x}{a}\right)^\alpha + \left(\frac{y}{b}\right)^\beta = 1$  where  $\alpha > 0$ ,  $\beta > 0$  given that density at any point  $p(xy)$  is  $k\sqrt{xy}$ .



- (e) Using the divergence theorem, evaluate the surface integral  $\iint_S (yz \, dy \, dz + zx \, dz \, dx + xy \, dy \, dx)$  where  $S : x^2 + y^2 + z^2 = 4$ .

### SECTION - C

Attempt any two parts from each question. All questions are compulsory.

3 (a) Trace the curve  $r^2 = a^2 \cos 2\theta$

(b) If  $u = \log \left( \frac{x^2 + y^2}{x + y} \right)$ , prove that

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 1$$

(c) If  $V = f(2x - 3y, 3y - 4z, 4z - 2x)$ , compute the value of  $6V_x + 4V_y + 3V_z$ .

(a) The temperature ' $T$ ' at any point  $(xyz)$  in space is  $T(xyz) = K xyz^2$  where  $K$  is constant. Find the highest temperature on the surface of the sphere  $x^2 + y^2 + z^2 = a^2$ .

(b) Verify the chain rule for Jacobians if  $x = u$ ,  $y = u \tan v$ ,  $z = w$ .



- (c) The time ' $T$ ' of a complete oscillation of a simple pendulum of length ' $L$ ' is governed by the equation

$$T = 2\pi \sqrt{\frac{L}{g}}, \quad g \text{ is constant, find the approximate}$$

error in the calculated value of  $T$  corresponding to an error of 2% in the value of  $L$ .

- 5 (a) Determine ' $b$ ' such that the system of homogeneous equation  $2x + y + 2z = 0$ ;  $x + y + 3z = 0$ ;  $4x + 3y + bz = 0$  has (i) Trivial solution (ii) Non-Trivial solution. Find the Non-Trivial solution using matrix method.

- (b) Verify Cayley-Hamilton theorem for the matrix

$$A = \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix} \text{ and hence find } A^{-1}.$$

- (c) Find the eigen value and corresponding eigen vectors of the matrix

$$I = \begin{pmatrix} -5 & 2 \\ 2 & -2 \end{pmatrix}$$

- 6 (a) Find the directional derivative of  $\nabla(\nabla f)$  at the point  $(1, -2, 1)$  in the direction of the normal to the surface  $xy^2z = 3x + z^2$  where  $f = 2x^3y^2z^4$ .



- (b) Using Green's theorem, find the area of the region in the first quadrant bounded by the curves

$$y = x, y = \frac{1}{x}, y = \frac{x}{4}$$

- (c) Prove that  $(y^2 - z^2 + 3yz)\hat{i} + (3xz + 2xy)\hat{j} + (3xy - 2xz + 2z)\hat{k}$  is both solenoidal and irrotational.

- 7 (a) Changing the order of integration of

$$\int_0^{\infty} \int_0^{\infty} e^{-xy} \sin nx \, dx \, dy$$

Show that  $\int_0^{\infty} \left( \frac{\sin nx}{x} \right) dx = \frac{\pi}{2}$ .

- (b) Determine the area bounded by the curves  $xy = 2$ ,  $4y = x^2$  and  $y = 4$ .
- (c) For a  $\beta$  function, show that  $\beta(p, q) = \beta(p+1, q) + \beta(p, q+1)$ .

