

(Following Paper ID and Roll No. to be filled in your Answer Book)

**PAPER ID : 9601**

Roll No.

--	--	--	--	--	--	--	--	--	--

## B. Tech.

(Semester-I) Theory Examination, 2012-13

### MATHEMATICS-I

*Time : 3 Hours]*

*[Total Marks : 100*

*Note :* Attempt questions from each Section as per instructions. The symbols have their usual meaning.

#### *Section-A*

Attempt *all* parts of this question. Each part carries 2 marks. 2×10=20

1. (a) Find  $y_n$ , if  $y = \frac{ax+b}{cx+d}$ .
- (b) Find all the asymptotes of the curve  $xy^2 = 4a^2(2a-x)$ .

(c) If  $z = xyf\left(\frac{x}{y}\right)$ , show that :

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 2z.$$

(d) Determine the point(s) where the function  $u = x^2 + y^2 + 6x + 12$  has a maximum or minimum.

(e) Evaluate  $\frac{\binom{8}{3}}{\binom{2}{3}}$ .

(f) Change the order of integration :

$$\int_0^a \int_0^{2\sqrt{ay}} f(x, y) dx dy + \int_0^a \int_0^{a-y} f(x, y) dx dy$$

(g) If  $\vec{A}$  and  $\vec{B}$  are irrotational, prove that  $\vec{A} \times \vec{B}$  is solenoidal.

(h) Find  $\int_C \hat{t} \cdot d\vec{r}$ , where  $\hat{t}$  is the unit tangent vector and  $C$  is the unit circle in the  $xy$ -plane about the origin.

(i) If  $A$  is a skew-Hermitian matrix, prove that  $(iA)$  is Hermitian matrix.

- (j) Find the sum and product of eigenvalues of the matrix :

$$\begin{bmatrix} 1 & 6 & 1 \\ 1 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

**Section-B**

Attempt any *three* parts of this question. Each part carries 10 marks. 10×3=30

2. (a) Find  $y_3$  when  $y = \sqrt{1+x^2} \cdot \sin x$  by Leibnitz theorem.

- (b) If  $u = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$ , then prove that :

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0.$$

- (c) Prove that :

$$\iiint \frac{dx dy dz}{\sqrt{a^2 - x^2 - y^2 - z^2}} = \frac{\pi^2 a^2}{8},$$

the integral being extended for all positive values of the variables for which the expression is real.

- (d) Apply Stoke's theorem to evaluate :

$$\oint_C [(x+y)dx + (2x-z)dy + (y+z)dz],$$

where  $C$  is the boundary of the triangle with vertices  $(2, 0, 0)$ ,  $(0, 3, 0)$  and  $(0, 0, 6)$ .

- (e) Find the square matrix  $A$  whose eigenvalues are 1, 2 and 3 and their corresponding eigenvectors are  $[1, 0, -1]^t$ ,  $[0, 1, 0]^t$  and  $[1, 0, 1]^t$  respectively.

### Section-C

Attempt *all* questions of this Section. Attempt any *two* parts from each question. Each question carries 10 marks.

$$10 \times 5 = 50$$

3. (a) If  $y = x^n \ln x$ , prove that  $xy_{n+1} = n!$ .  
(b) Prove that :

$$f\left(\frac{x^2}{1+x}\right) = f(x) - \frac{x}{1+x} f'(x) + \frac{1}{2!} \frac{x^2}{(1+x)^2} f''(x) - \dots$$

- (c) Trace the curve  $y^2(a-x) = x^2(a+x)$ .

4. (a) If  $u = x_1 + x_2 + x_3 + x_4$ ,  $uv = x_2 + x_3 + x_4$ ,  
 $uvw = x_3 + x_4$  and  $uvwt = x_4$ , find :

$$\frac{\partial(x_1, x_2, x_3, x_4)}{\partial(u, v, w, t)}$$

- (b) What error in the common logarithm of a number will be produced by an error of 1% in the number ?
- (c) If  $x^x \cdot y^y \cdot z^z = C$ , show at  $x = y = z$  :

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{-1}{x \ln(ex)}$$

5. (a) Evaluate the integral :

$$\int_0^{\infty} \int_x^{\infty} \frac{e^{-y}}{y} dy dx.$$

- (b) Find, by double integration, the area of the region enclosed by the curves  $x^2 + y^2 = a^2$ ,  
 $x + y = a$  in the first quadrant.
- (c) Show that :

$$\int_0^1 \frac{dx}{\sqrt{1-x^n}} = \frac{\sqrt{\pi} \cdot \sqrt{\frac{1}{n}}}{n \cdot \sqrt{\left(\frac{1}{n} + \frac{1}{2}\right)}}$$

6. (a) If  $\phi(x, y) = \frac{1}{2} \ln(x^2 + y^2)$ , show that :

$$\text{grad } \phi = \frac{\vec{r} - (\hat{k} \cdot \vec{r}) \hat{k}}{\{\vec{r} - (\hat{k} \cdot \vec{r}) \hat{k}\} \cdot \{\vec{r} - (\hat{k} \cdot \vec{r}) \hat{k}\}}$$

- (b) If  $u = x + y + z$ ,  $v = x^2 + y^2 + z^2$  and  $w = xy + yz + zx$ , show that  $\text{grad } u$ ,  $\text{grad } v$  and  $\text{grad } w$  are coplanar.
- (c) Consider a vector field :

$$\vec{F} = (x^2 - y^2 + x) \hat{i} - (2xy + y) \hat{j}.$$

Show that the field is irrotational and find its scalar potential. Hence evaluate the line integral from (1, 2) to (2, 1).

7. (a) Find the inverse of the matrix :

$$\begin{bmatrix} i & -1 & 2i \\ 2 & 0 & 2 \\ -1 & 0 & 1 \end{bmatrix}$$

by employing elementary transformations.

- (b) Find the value of  $\lambda$  such that the following equations have unique solution :

$$\lambda x + 2y - 2z = 1$$

$$4x + 2\lambda y - z = 2$$

$$6x + 6y + \lambda z = 3.$$

- (c) Examine the linear dependence of the vectors  $[1, -1, 1]$ ,  $[2, 1, 1]$  and  $[3, 0, 2]$ .  
If dependent, find the relation between them.