

Printed pages:3
Paper ID: 9 0 3 8

Roll No.

--	--	--	--	--	--	--	--	--	--

Sub Code: EAS-301

B-TECH
(SEM. III) THEORY EXAMINATION 2017-18
MATHEMATICS-III

Time: 3 Hours

[Total Marks: 100]

Note: 1. Attempt all Sections. If require any missing data; then choose suitably.

SECTION-A

1. Attempt all questions in brief. 2x10 = 20

- (a) State the necessary and sufficient conditions for $f(z)$ to be analytic.
- (b) If the first four moments of a distribution about the value 5 are equal to -4, 22, -117 and 560. Determine the corresponding moments about the mean.
- (c) The regression equations calculated from a given set of observation for two random variables are $x = -0.4y + 6.4$ and $y = -0.6x + 4.6$ calculate \bar{x}, \bar{y} .
- (d) Define poisson distribution.
- (e) Define the harmonic function.
- (f) State Simpson's 3/8 rule.
- (g) Find Z-transformation of $f(k) = \begin{cases} 1, & k = 0 \\ 0, & k \neq 0 \end{cases}$
- (h) Explain the false position method.
- (i) If $f(z)$ is analytic function of z show that $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)|f(z)|^2 = 4|f'(z)|^2$.
- (j) Define the Z - transform of the sequence $\{f_k\}$.

SECTION-B

2. Attempt any three parts from this section. (10×3=30)

- a. Using the Z-transform solve the following difference equations:

$$y_{k+2} + 4y_{k+1} + 3y_k = 3^k \text{ given } y_{(0)} = 0, y_{(1)} = 1.$$

- b. Using Cauchy Integral formula evaluate the following integral :

$$\int_C \frac{\sin \pi z^2 + \cos \pi z^2}{(z-1)(z-2)} dz \text{ where } C \text{ is } |Z| = 3$$

- c. Find the root of equation using Regula-Falsy Method correct up to four decimal places,

$$x - e^{-x} = 0 .$$

- d. Find the rank correlation coefficient from the following data :

X	74	75	78	72	78	77	79	81	79	76	72	71
Y	47	44	40	48	49	45	46	42	42	39	46	40

- e. Solve the following initial value problem

$$\frac{dy}{dx} = -2xy^2, y(0) = 1$$

With $h=0.2$ on interval $(0, 0.6)$ using fourth order Runge-Kutta method. Compare with exact solution.

SECTION-C

- 3. Attempt any one part of the following. 10x1 = 10**

(a) Expand $\frac{(z-2)(z+2)}{(z+1)(z+4)}$ in the regions (a) $|z| < 1$ (b) $1 < |z| < 4$.

(b) Evaluate the integral by contour integration $\int_0^{2\pi} \frac{\cos 3\theta}{5 - 4\cos\theta} d\theta$.

- 4. Attempt any one part of the following. 10x1 = 10**

(a) Find inverse Z-transformation of $\frac{z^2}{(z-1)(z-3)}$.

(b) Find the Fourier cosine transformation of $\frac{1}{1+x^2}$.

- 5. Attempt any one part of the following. 10x1 = 10**

- (a) Solve the following system of linear equations by Gauss-Seidel iteration Method:

$$x + y + 54z = 110; 27x + 6y - z = 85; 6x + 15y + 2z = 72$$

(b) Evaluate $\int_0^1 \frac{dx}{1+x^2}$ using Simpson's 1/3 rule, Simpson's 3/8 rule taking $h=1/4$?

- 6. Attempt any one part of the following. 10x1 = 10**

- (a) Fit a least square curve $y = ax^b$ to the given data

x	1	2	3	4	5	6
y	2.98	4.26	5.21	6.1	6.8	7.5

- (b) Use Newton's divided difference formula to find the interpolating polynomial and hence evaluate $y(9.5)$ from the given data :

x	7	8	9	10
y	3	1	1	9

- 7. Attempt any one part of the following. 10x1 = 10**

- (a) A river is 80 meter wide. The depth y of the river at a distance x from one bank is given by the following table:

x:	0	10	20	30	40	50	60	70	80
y:	0	4	7	9	12	15	14	8	3

Find the appropriate area of cross section of the river by using suitable Integration Formula.

- (b) Using Newton's Raphson Method solve the equation $\cos x - xe^x = 0$ correct to four decimal places.