



NAS-301

(Following Paper ID and Roll No. to be filled in your Answer Book)

PAPER ID : 199320

Roll No.

--	--	--	--	--	--	--	--	--	--

B. Tech.(SEM. III) (ODD SEM.) THEORY
EXAMINATION, 2014-15**MATHEMATICS-III**

Time : 3 Hours]

[Total Marks : 100

Note : Attempt All Questions. All Questions carry equal marks

1 Attempt any four parts of the following : 5×4=20

- (a) State Cauchy-Riemann theorem for an analytic function.
Test the analyticity of the following Function :

$$f(z) = \frac{(x^3 - y^3) + i(x^3 + y^3)}{x^2 + y^2}, \text{ if } z \neq 0 \text{ and}$$

$$= 0 \text{ if } z = 0$$

- (b) State Cauchy- integral theorem for an analytic function.
Verify this theorem by integrating the function $z^3 + iz$ along the boundary of the rectangle with vertices $+1, -1, i, -i$.
- (c) Show that the function $u = \frac{1}{2} \log(x^2 + y^2)$ is harmonic.

Find the harmonic conjugate of u.

- (d) Evaluate the integral $\int \frac{e^{2z}}{(z+1)^5} dz$, around the boundary

of the circle $|z| = 2$.

- (e) Find the Taylor series expansion of the function $\tan^{-1} z$ about the point $z = \pi/4$.

- (f) Evaluate the integral $\int_0^\pi \frac{\cos^2 3\theta}{5 - 4 \cos 2\theta} d\theta$.

2 Attempt any two parts of the following : 10×2=20

- (a) Find the Fourier transform of the following function $f(x) = 1 - x^2$, if $|x| \leq 1$ and $f(x) = 0$, if $|x| > 1$
- (b) Using Z — transform solve the following difference equation

$$Y_{n+2} - (2 \cos \alpha) Y_{n+1} + Y_n = 7^n \text{ with the conditions that } Y_0 = 5, Y_1 = 1.$$

- (c) State the convolution theorem for Fourier transform.
Prove that the Fourier transform of the convolution of the two functions equal to the product of their Fourier transforms.

3 Attempt any two parts of the following : 10×2=20

- (a) Define skewness and kurtosis of a distribution. The first four moments of a distribution are 0, 2.5, 0.7, and 18.71. Find the coefficient of skewness and kurtosis.
- (b) Fit a second degree parabola to the following data :

x	1	2	3	4	5	6	7	8	9
y	2	6	7	8	10	11	8	13	5

- (c) Define coefficient of correlation and regression.
If θ is the acute angle between the two lines of regression then prove that

$$\tan \theta = \frac{1-r^2}{r} \frac{\sigma_x \sigma_y}{\sigma_x^2 + \sigma_y^2}$$

where r , σ_x , σ_y have their usual meanings. Give the significance of the formula when $r = 0$ and $r = \pm 1$.

- 4 Attempt any two parts of the following : **10×2=20**

- (a) Derive Newton – Raphson's method to find a root of the equation $f(x) = 0$. Prove that this method has quadratic convergence.
(b) Apply Newton's divided difference method to obtain an interpolatory polynomial for the following data.

x	3	5	7	9	11	13
$f(x)$	31	51	17	19	90	110

- (c) Obtain Lagrange's Interpolatory for the following data :

x	1	3	5	7	10
$f(x)$	13	31	25	37	101

Find the values of $f(4)$ and $f(8.5)$.

- 5 Attempt any two parts of the following : **10×2=20**

- (a) Solve the following system of linear equations using Gauss-Seidel method
 $10x + 3y + 7z = 41$,
 $3x + 20y + 17z = 101$,
 $x + 19y + 23z = 201$,
 perform three iterations.
 (b) State Simpson's three-eighth rule. Using this rule evaluate the following integral

$$\int_0^6 \frac{x}{1+x^5} dx$$

- (c) State Runge-Kutta method of fourth order. Using this method find the values of $y(0.2)$, $y(0.4)$ and $y(0.6)$ for the following initial value problem

$$\frac{dy}{dx} = x^3 - y^3 \text{ with condition that } y(0) = 1$$

UPTUonline.com