

Printed Pages : 3



NCH401

(Following Paper ID and Roll No. to be filled in your Answer Book)

PAPER ID : 151409

Roll No.

--	--	--	--	--	--	--	--	--	--

B. Tech.

(SEM. IV) THEORY EXAMINATION, 2014-15

THERMODYNAMICS - I

Time : 2 Hours]

[Total Marks : 50

Note: Attempt all questions.**1** Attempt **any four** parts of the following: **3×4=12**

- (a) Explain the concept of First Law of Thermodynamics and law of conservation of energy.
- (b) Given a thermodynamic system, conduct an analysis using the Second Law of Thermodynamics.
- (c) Discuss the virial expressions for compressibility factor Z .
- (d) Describe triple point and critical point for a pure substance.
- (e) Define ideal solution. What is Lewis/Randall rule?
- (f) What are the effects of the temperature on the heat of reaction ?

151409]

1

[Contd...

2 Attempt **any two** parts of the followings: **6×2=12**

- (a) What is change in entropy when 0.7 m^3 of CO_2 and 0.3 m^3 of N_2 each at 1 bar and 25°C blend to form a gas mixture at the same condition? Assume ideal gases.
- (b) A heat engine which rejects waste heat to a sink at 310 K has a thermal efficiency of 36% and a second law efficiency of 60%. Determine the temperature of the source that supplies heat to this heat engine.
- (c) Show T xy and P xy diagram for binary system of immiscible liquids.

3 Attempt **any two** parts of the followings: **7×2=14**

- (a) One cubic meter of an ideal gas at 600 K and 1000 kPa expands to four times its initial volume by a mechanically reversible adiabatic process. Calculate the temperature, pressure and work done by the gas. Take $C_p = 2.5/\text{mol K}$.
- (b) A 30 kg steel casting ($C_p = 0.5 \text{ kJ kg}^{-1} \text{ K}^{-1}$) at a temperature of 723 K is quenched in 150 kg of oil ($C_p = 2.55 \text{ kJ kg}^{-1} \text{ K}^{-1}$) at 298 K. If there is no heat losses what is change in' entropy of oil?
- (c) Write short notes on:
 - (i) Enthalpy concentration diagram
 - (ii) Gibbs - Duhem equation.

4 Attempt **any two** parts of the followings: **6×2=12**

- (a) A Carnot engine operates between temperature levels 600 K and 300 K. It drives a Carnot refrigerator, which provides cooling at 250 K and discards heat at 300K. Determine a numerical value for the ratio of heat extracted by the refrigerator to the heat delivered to the engine.
- (b) Derive Fundamental property relation.
- (c) A closed nonreactive system contains species 1 and 2 in vapour/liquid equilibrium. Species 2 is very light gas, essentially insoluble in the liquid phase. The vapour phase contains both species 1 and 2. Some additional moles of species 2 are added to the system, which is then restored to its initial T and P. As a result of process, does the total number of moles of liquid increase, decrease or remains unchanged? Give explanation in support of your answer.
-